

DESCRIPTION

Target:	F3
Target aliases:	Coagulation Factor III, TF, CD142
Fc isotype:	IgG2a
Membrane proteome specificity:	Monospecific for 6,000 membrane proteins tested
Species reactivity:	Human (Others untested)
Extracellular epitope:	Yes
Fc modifications:	C-terminal Avitag ¹ , disabled Fc- γ receptor binding ²
Source:	Recombinant CHO expression; purified by Protein A chromatography
Formulation:	Endotoxin Free PBS pH 7.4, sterile-filtered
Concentration:	1 mg/ml


1. A peptide tag that can be biotinylated in vitro using the biotin ligase enzyme (BirA).
2. Mutated Fc- γ receptor binding site to minimize non-specific antibody binding.

F3 TARGET INFORMATION


F3 is a cell surface glycoprotein with extracellular, transmembrane, and cytoplasmic domains that binds coagulation factor VII, initiating the coagulation protease cascade. F3 is involved with HIV-associated coagulopathy, and platelet-dependent monocyte expression of this gene is associated with COVID-19 severity and mortality. (NCBI Gene: 2152, UniProtKB/Swiss-Prot: P13726) Other names: Coagulation Factor III, TF, CD142

SHIPPING AND STORAGE

Shipping:	Shipped at ambient temperature. Store at 4°C.
Stability & Storage:	Stable for 12 months from date of receipt when stored at 4°C. Avoid repeated freeze-thaw cycles.


VALIDATION DATA
Membrane Proteome Specificity


Applications	Conditions	Recommended concentration
Flow Cytometry, Extracellular	Live, Unpermeabilized	1 µg/ml



HEK-293F cells transiently transfected with F3 were stained with F3 Monoclonal Antibody (CSB0010) (green) or isotype control antibody (gray), followed by AlexaFluor 647- conjugated anti- IgG secondary antibody. HEK-293F cells transiently transfected with an empty control vector were also stained with F3 Monoclonal Antibody (CSB0010) (blue).

Applications	Conditions	Recommended concentration
Immunofluorescence, Extracellular	Fixed 4% paraformaldehyde	1 µg/ml



(Left) COS-7 cells transiently transfected F3 were stained with F3 Monoclonal Antibody (CSB0010) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) COS-7 cells transiently transfected with an empty control vector stained with F3 Monoclonal Antibody. (Bottom right) Isotype control: COS-7 cells transfected with F3 and stained with control MAb.

Applications


Immunofluorescence, Intracellular

Conditions

Fixed 4% paraformaldehyde,
Permeabilized 0.1% Triton X-100

Recommended concentration

1 µg/ml



(Left) COS-7 cells transiently transfected F3 were permeabilized and stained with F3 Monoclonal Antibody (CSB0010) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) COS-7 cells transiently transfected with an empty control vector stained with F3 Monoclonal Antibody. (Bottom right) Isotype control: COS-7 cells transfected with {{target}} and stained with control MAb.