

DESCRIPTION

Target:	MET
Target aliases:	DFNB97, RCCP2, HGFR, SF Receptor, AUTS9, C-Met
Fc isotype:	IgG2a
Membrane proteome specificity:	Monospecific for 6,000 membrane proteins tested
Species reactivity:	Human (Others untested)
Extracellular epitope:	Yes
Fc modifications:	C-terminal Avitag ¹ , disabled Fc- γ receptor binding ²
Source:	Recombinant CHO expression; purified by Protein A chromatography
Formulation:	Endotoxin Free PBS pH 7.4, sterile-filtered
Concentration:	1 mg/ml

1. A peptide tag that can be biotinylated in vitro using the biotin ligase enzyme (BirA).
2. Mutated Fc- γ receptor binding site to minimize non-specific antibody binding.

MET TARGET INFORMATION


MET is a single-pass transmembrane protein and receptor tyrosine kinase with a secreted isoform. MET binds hepatocyte growth factor and is involved in embryonic development and cellular migration, invasion, and survival. Mutations in MET are associated with various types of cancer. (NCBI Gene: 4233, UniProtKB/Swiss-Prot: P08581) Other names: DFNB97, RCCP2, HGFR, SF Receptor, AUTS9, C-Met

SHIPPING AND STORAGE


Shipping:	Shipped at ambient temperature. Store at 4°C.
Stability & Storage:	Stable for 12 months from date of receipt when stored at 4°C. Avoid repeated freeze-thaw cycles.

VALIDATION DATA

Membrane Proteome Specificity




Applications	Conditions	Recommended concentration
Flow Cytometry, Extracellular	Live, Unpermeabilized	1 µg/ml



JS-1 cells transiently transfected with MET were stained with MET Monoclonal Antibody (CSB0070) (green) or isotype control antibody (gray), followed by AlexaFluor 647- conjugated anti- IgG secondary antibody. JS-1 cells transiently transfected with an empty control vector were also stained with MET Monoclonal Antibody (CSB0070) (blue).

Applications	Conditions	Recommended concentration
Immunofluorescence, Extracellular	Fixed 4% paraformaldehyde	1 µg/ml



(Left) JS-1 cells transiently transfected MET were stained with MET Monoclonal Antibody (CSB0070) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) JS-1 cells transiently transfected with an empty control vector stained with MET Monoclonal Antibody. (Bottom right) Isotype control: JS-1 cells transfected with MET and stained with control MAb.

Applications


Immunofluorescence, Intracellular

Conditions

Fixed 4% paraformaldehyde,
Permeabilized 0.1% Triton X-100

Recommended concentration

1 µg/ml



(Left) JS-1 cells transiently transfected MET were permeabilized and stained with MET Monoclonal Antibody (CSB0070) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) JS-1 cells transiently transfected with an empty control vector stained with MET Monoclonal Antibody. (Bottom right) Isotype control: JS-1 cells transfected with {{target}} and stained with control MAb.