

DESCRIPTION

Target:	CD1D
Target aliases:	R3G1, CD1A, R3
Fc isotype:	IgG2a
Membrane proteome specificity:	Monospecific for 6,000 membrane proteins tested
Species reactivity:	Human (Others untested)
Extracellular epitope:	Yes
Fc modifications:	C-terminal Avitag ¹ , disabled Fc- γ receptor binding ²
Source:	Recombinant CHO expression; purified by Protein A chromatography
Formulation:	Endotoxin Free PBS pH 7.4, sterile-filtered
Concentration:	1 mg/ml

1. A peptide tag that can be biotinylated in vitro using the biotin ligase enzyme (BirA).
2. Mutated Fc- γ receptor binding site to minimize non-specific antibody binding.

CD1D TARGET INFORMATION


CD1D is a single-pass transmembrane protein that is part of the CD1 family of transmembrane glycoproteins. CD1D, like other members of the CD1 family, binds and presents self and microbial glycolipids to T-cells. (NCBI Gene: 912, UniProtKB/Swiss-Prot: P15813) Other names: R3G1, CD1A, R3

SHIPPING AND STORAGE

Shipping:	Shipped at ambient temperature. Store at 4°C.
Stability & Storage:	Stable for 12 months from date of receipt when stored at 4°C. Avoid repeated freeze-thaw cycles.

VALIDATION DATA

Membrane Proteome Specificity



The specificity of CD1D Monoclonal Antibody (CSB0094) was tested on the Membrane Proteome Array™ and shown to be specific for CD1D.

The Membrane Proteome Array™ contains 6,000 different human membrane proteins, each expressed in unfixed human cells to ensure native conformation and post-translational modifications. The Membrane Proteome Array™ represents the industry standard for determining the binding specificity of antibodies and other protein ligands.

Applications	Conditions	Recommended concentration
Immunofluorescence, Extracellular	Fixed 4% paraformaldehyde	1 µg/ml
mAb + target mAb + control Control mAb + target © Cell Surface Bio (Left) COS-7 cells transiently transfected CD1D were stained with CD1D Monoclonal Antibody (CSB0094) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) COS-7 cells transiently transfected with an empty control vector stained with CD1D Monoclonal Antibody. (Bottom right) Isotype control: COS-7 cells transfected with CD1D and stained with control MAb.		