

DESCRIPTION

Target:	CD4
Target aliases:	CD4mut, IMD79, OKT4D, T-Cell Surface Antigen T4/Leu-3
Fc isotype:	IgG2a
Membrane proteome specificity:	Monospecific for 6,000 membrane proteins tested
Species reactivity:	Human (Others untested)
Extracellular epitope:	Yes
Fc modifications:	C-terminal Avitag ¹ , disabled Fc- γ receptor binding ²
Source:	Recombinant CHO expression; purified by Protein A chromatography
Formulation:	Endotoxin Free PBS pH 7.4, sterile-filtered
Concentration:	1 mg/ml


1. A peptide tag that can be biotinylated in vitro using the biotin ligase enzyme (BirA).
2. Mutated Fc- γ receptor binding site to minimize non-specific antibody binding.

CD4 TARGET INFORMATION

Cluster of differentiation 4 (CD4) is a type I transmembrane glycoprotein belonging to the immunoglobulin superfamily. It is primarily expressed on the surface of T helper lymphocytes, monocytes, macrophages, and dendritic cells, where it functions as a co-receptor that enhances T cell receptor signaling through interaction with major histocompatibility complex class II (MHC II) molecules. The extracellular region contains four immunoglobulin-like domains, while the cytoplasmic tail associates with the tyrosine kinase LCK to facilitate intracellular signaling. CD4 also serves as a receptor for the HIV-1 envelope glycoprotein gp120, enabling viral entry into host cells. (NCBI Gene: 920, UniProtKB/Swiss-Prot: P01730) Other names: CD4mut, IMD79, OKT4D, T-Cell Surface Antigen T4/Leu-3

SHIPPING AND STORAGE


Shipping:	Shipped at ambient temperature. Store at 4°C.
Stability & Storage:	Stable for 12 months from date of receipt when stored at 4°C. Avoid repeated freeze-thaw cycles.

VALIDATION DATA
Membrane Proteome Specificity


The specificity of CD4 Monoclonal Antibody (CSB0118) was tested on the Membrane Proteome Array™ and shown to be specific for CD4.


The Membrane Proteome Array™ contains 6,000 different human membrane proteins, each expressed in unfixed human cells to ensure native conformation and post-translational modifications. The Membrane Proteome Array™ represents the industry standard for determining the binding specificity of antibodies and other protein ligands.

Applications	Conditions	Recommended concentration
Flow Cytometry, Extracellular	Live, Unpermeabilized	1 µg/ml






HEK-293F cells transiently transfected with CD4 were stained with CD4 Monoclonal Antibody (CSB0118)(green) or isotype control antibody (gray), followed by AlexaFluor 647- conjugated anti- IgG secondary antibody. HEK-293F cells transiently transfected with an empty control vector were also stained with CD4 Monoclonal Antibody (CSB0118)(blue).

Applications	Conditions	Recommended concentration
Immunofluorescence, Extracellular	Fixed 4% paraformaldehyde	1 µg/ml



(Left) COS-7 cells transiently transfected CD4 were stained with CD4 Monoclonal Antibody (CSB0118) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) COS-7 cells transiently transfected with an empty control vector stained with CD4 Monoclonal Antibody. (Bottom right) Isotype control: COS-7 cells transfected with CD4 and stained with control MAb.

Applications	Conditions	Recommended concentration
Immunofluorescence, Intracellular	Fixed 4% paraformaldehyde, Permeabilized 0.1% Triton X-100	1 µg/ml
mAb + target  © Cell Surface Bio mAb + control  Control mAb + target  (Left) COS-7 cells transiently transfected CD4 were permeabilized and stained with CD4 Monoclonal Antibody (CSB0118) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) COS-7 cells transiently transfected with an empty control vector stained with CD4 Monoclonal Antibody. (Bottom right) Isotype control: COS-7 cells transfected with {{target}} and stained with control MAb.		