

DESCRIPTION

Target:	CNR1
Target aliases:	CB1, CANN6, CB-R, CB1K5, CB1A, CNR, CB1R
Fc isotype:	IgG2a
Membrane proteome specificity:	Monospecific for 6,000 membrane proteins tested
Species reactivity:	Human (Others untested)
Extracellular epitope:	Yes
Fc modifications:	C-terminal Avitag ¹ , disabled Fc- γ receptor binding ²
Source:	Recombinant CHO expression; purified by Protein A chromatography
Formulation:	Endotoxin Free PBS pH 7.4, sterile-filtered
Concentration:	1 mg/ml

1. A peptide tag that can be biotinylated in vitro using the biotin ligase enzyme (BirA).
2. Mutated Fc- γ receptor binding site to minimize non-specific antibody binding.

CNR1 TARGET INFORMATION


CNR1 is a multi-pass transmembrane protein and cannabinoid receptor. Cannabinoid receptors are part of the G-protein coupled receptor family and mediate the central nervous system effects of marijuana. (NCBI Gene: 1268, UniProtKB/Swiss-Prot: P21554) Other names: CB1, CANN6, CB-R, CB1K5, CB1A, CNR, CB1R

SHIPPING AND STORAGE

Shipping:	Shipped at ambient temperature. Store at 4°C.
Stability & Storage:	Stable for 12 months from date of receipt when stored at 4°C. Avoid repeated freeze-thaw cycles.

VALIDATION DATA

Membrane Proteome Specificity




Applications	Conditions	Recommended concentration
Flow Cytometry, Extracellular	Live, Unpermeabilized	1 µg/ml

HEK-293F cells transiently transfected with CNR1 were stained with CNR1 Monoclonal Antibody (CSB0129) (green) or isotype control antibody (gray), followed by AlexaFluor 647- conjugated anti- IgG secondary antibody. HEK-293F cells transiently transfected with an empty control vector were also stained with CNR1 Monoclonal Antibody (CSB0129) (blue).

Applications	Conditions	Recommended concentration
Immunofluorescence, Extracellular	Fixed 4% paraformaldehyde	1 µg/ml

(Left) JS-1 cells transiently transfected CNR1 were stained with CNR1 Monoclonal Antibody (CSB0129) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) JS-1 cells transiently transfected with an empty control vector stained with CNR1 Monoclonal Antibody. (Bottom right) Isotype control: JS-1 cells transfected with CNR1 and stained with control MAb.

Applications	Conditions	Recommended concentration
Immunofluorescence, Intracellular	Fixed 4% paraformaldehyde, Permeabilized 0.1% Triton X-100	1 µg/ml



(Left) JS-1 cells transiently transfected CNR1 were permeabilized and stained with CNR1 Monoclonal Antibody (CSB0129) followed by AlexaFluor 647 anti- IgG secondary antibody (red) and DAPI (blue). (Top right) JS-1 cells transiently transfected with an empty control vector stained with CNR1 Monoclonal Antibody. (Bottom right) Isotype control: JS-1 cells transfected with {{target}} and stained with control MAb.